

Justifying the exotic Θ^+ pentaquark

Dmitri Diakonov

Petersburg Nuclear Physics Institute, Gatchina 188300, St. Petersburg, Russia

Abstract

The existence of a light $S = +1$ baryon resonance follows from Quantum Field Theory applied to baryons. This is illustrated in the Skyrme model (where Θ^+ exists but is too strong) and in a new mean field approach where Θ^+ arises as a consequence of three known resonances: $\Lambda(1405)$, $N(1440)$ and $N(1535)$.

Key words: baryon resonances, exotic resonances, Skyrme model, pentaquark

PACS: 12.39.Ki 12.38.Lg 12.39.Mk 13.30.Eg

After strong signals of the exotic $S = +1$ baryon have been announced in the Fall of 2002 from two independent searches [1,2] initiated by a theoretical prediction of a *light and narrow* exotic resonance [3], there has been much confusion in the field, both on the experimental and theoretical sides. The anomalously small width is the key to that confusion. If $\Gamma_\Theta \leq 1$ MeV [4] it means that the (ΘKN) coupling is an order of magnitude less than the typical pseudoscalar couplings of normal mainly $3Q$ baryons, and one expects that the vector coupling (ΘK^*N) is similarly strongly suppressed. The Θ^+ production cross sections are then very small and it indeed becomes a challenge to reveal it experimentally.

Theoretically, both the light mass and a small width are uncomprehensible from the naïve point of view on hadrons, which assumes one adds up ≈ 350 MeV per constituent quark, plus 150 MeV for strangeness. With that arithmetics, one expects a pentaquark $uudd\bar{s}$ state at ≥ 1900 MeV (400 MeV heavier than in reality!) and, of course, there is no way to avoid a large width from a fall-apart decay.

There have been strong warnings in the past that the constituent quark picture is an oversimplification, the “spin crisis” and the anomalously huge nucleon σ -term being examples of such warnings [5]. However it is the light and narrow Θ^+ that is really a shock to the simplistic view. Not only is Θ^+ exotic but its width is unprecedentedly small for a strongly decaying resonance.

Lacking in the traditional picture is Quantum Field Theory saying that all baryons are in fact quantum-mechanical superpositions of $3Q, 5Q, 7Q, \dots$ states, and Spontaneous

Breaking of Chiral Symmetry, saying that constituent quarks *have* to interact strongly at least with the pseudoscalar meson fields. A very rough, very approximate model of baryons that accommodates, however, this physics is the Skyrme model. There is a prejudice that the Skyrme model is somehow opposite or perpendicular to quark models. In fact the Skyrme model is in accordance with quarks [6] but it is in one way better as it allows baryons to be made of N_c quarks, plus an indefinite number of $Q\bar{Q}$ pairs.

The question whether Θ^+ exists in the Skyrme model has been studied, at large N_c , in Ref. [7] in the Callan–Klebanov approach, with the conclusion that there are “no $S=+1$ kaon bound states or resonances in the spectrum”. This is a misunderstanding: the KN cross section in the $T=0, L=1$ wave, computed from the phase shifts in the Skyrme model [7] exhibits a strong resonance around 1500 MeV, see Fig. 1 (the existing data for the full cross section are also shown there). A more precise way to formulate it is that the KN scattering amplitude has a pole at $m_{\text{res}} - i\frac{\Gamma}{2} = 1449 - i44$ MeV [6], for the standard




Fig. 1. Skyrme model predicts a strong Θ resonance.




Fig. 2. Ground-state baryon $N(940, 1/2^+)$.

parameters of the Skyrme model. Therefore, the Skyrme model *does* predict a strong exotic resonance, and there is no way to get rid of it as its origin is very general. However, as explained in detail in Ref. [6], the Skyrme model grossly overestimates the Θ^+ width. In realistic settings it becomes very narrow, and that is why it is so hard to detect it.

There is a nice way to understand Θ^+ in simple terms and see that it is unavoidable [8]. If the number of quark colors $N_c=3$ is considered to be a large number, the N_c quarks in a baryon can be viewed, according to Witten, as bound by a mean field. Any reasonable *Ansatz* for the mean field breaks symmetry between u, d quarks on the one hand and s quarks on the other. The (approximate) $SU(3)$ symmetry is restored when one considers rotations of the mean field in flavor space: that produces $SU(3)$ baryon multiplets. The splittings between multiplet centers are $\mathcal{O}(1/N_c)$, and the splittings inside multiplets are $\mathcal{O}(m_s)$. The ground-state baryon – the nucleon – is obtained by filling in all negative-energy one-particle states in the mean field, and adding one filled shell with positive energy for u, d quarks, see Fig. 2. The s -quark shells are characterized by J^P whereas the u, d -quark shells are characterized by K^P where $\mathbf{K}=\mathbf{J}+\mathbf{T}$. The highest shell filled by N_c quarks in antisymmetric state in color must have $J^P = 1/2^+$ for s quarks, and $K^P = 0^+$ for u, d quarks. The lowest baryon multiplets $(\mathbf{8}, 1/2^+)$ and $(\mathbf{10}, 3/2^+)$ are obtained from quantizing the rotations of this filling scheme in flavor and ordinary spaces.

The lowest baryon resonances that are not rotational excitations of the ground-state baryon are $\Lambda(1405, 1/2^-)$ and $N(1440, 1/2^+)$. They can be obtained as one-quark exci-

tations in the mean field: $\Lambda(1405)$ as the $1/2^-$ excitation of the s quark (Fig. 3) and $N(1440)$ as a 0^+ (or 1^+) excitation of u, d quarks (Fig. 4). The existence of the excited $1/2^-$ level for s quarks implies that it can be also excited by s quark from the highest filled s -quark shell: such particle-hole excitation can be identified with the $N(1535, 1/2^-)$ resonance, see Fig. 3. At $N_c=3$ it is mainly a pentaquark baryon $u(d)uds\bar{s}$, which explains its large decay branching into ηN , a long-time mystery.




Fig. 3. One-quark excitations corresponding to $\Lambda(1405, 1/2^-)$ and $N(1535, 1/2^-)$.




Fig. 4. One-quark excitations corresponding to $N(1440, 1/2^+)$ and $\Theta^+(1/2^+)$.

The existence of the excited (Roper) 0^+ level for u, d quarks implies that it can be also excited from the s -quark shell, forming an exotic pentaquark $\Theta^+ = uud\bar{s}$, see Fig. 4. Since the positions of all levels involved are already fixed from the masses of the three resonances, we estimate the Θ^+ mass as $m_\Theta \approx 1440 + 1535 - 1405 = 1570$ MeV, with the uncertainty of few tens MeV as the resonance masses are not precisely known. Thus, in the mean field picture the exotic pentaquark is a consequence of the three well-known resonances and is light [8]. One should not expect a qualitative change in the levels as one goes from large N_c , where the mean field is exact, to the real-world $N_c=3$.

In this interpretation, Θ^+ is a Gamov–Teller-type resonance long known in nuclear physics, where a neutron from a filled shell can be put on an excited proton level. All four excitations shown in Figs. 3,4 entail their own rotational bands of $SU(3)$ multiplets.

Finally, I would like to remark that with all couplings of Θ^+ to ordinary baryons being small, a promising way of detecting it is in interference with the production of known resonances, since the interference cross sections are linear (and not quadratic) in the small coupling constants [9].

I acknowledge many helpful discussions with Victor Petrov and Klaus Goeke.

References

- [1] T. Nakano [LEPS Collaboration], Talk at the PANIC 2002 (Oct. 3, 2002, Osaka); T. Nakano *et al.*, Phys. Rev. Lett. **91**, 012002 (2003)
- [2] V.A. Shebanov [DIANA Collaboration], Talk at the Session of the Nuclear Physics Division of the RAS (Dec. 3, 2002, Moscow); V.V. Barmin *et al.*, Phys. Atom. Nucl. **66**, 1715 (2003)
- [3] D. Diakonov, V. Petrov and M. Polyakov, Z. Phys. **A359**, 305 (1997); arXiv:hep-ph/0404212
- [4] V.V. Barmin *et al.* [DIANA Collaboration], Phys. Atom. Nucl. **70**, 35 (2007), arXiv:hep-ex/0603017
- [5] D. Diakonov, Talk at the APS Meeting, Denver, Colorado, 1-4 May 2004, arXiv:hep-ph/0406043
- [6] D. Diakonov and V. Petrov, to be published by World Scientific in *Multifaceted Skyrmions*, G. Brown and M. Rho (eds.), arXiv:0812.1212
- [7] N. Itzhaki, I.R. Klebanov, P. Ouyang and L. Rastelli, Nucl. Phys. **B684**, 264 (2004)
- [8] D. Diakonov, arXiv:0812.3418
- [9] M. Amarian, D. Diakonov and M. Polyakov, Phys. Rev. **D78**, 074003 (2008), arXiv:hep-ph/0612150